

A TOY MODEL FOR GAMMA-RAY BURSTS IN TYPE IB/C SUPERNOVAE

WEI-HUA LEI, DING-XIONG WANG¹, AND REN-YI MA

Department of Physics, Huazhong University of Science and Technology, Wuhan, 430074, People's Republic of China

dxwang@hust.edu.cn

ABSTRACT

A toy model for gamma-ray burst-supernovae (GRB-SN) is discussed by considering the coexistence of baryon poor outflows from black holes (BHs) and a powerful spin-connection to surrounding disk, giving rise to consistent calorimetry as described by van Putten (2001a) in a variant of the Blandford-Znajek process (BZ, 1977). In this model the half-opening angle of the magnetic flux tube on the horizon is determined by the mapping relation between the angular coordinate on the BH horizon and the radial coordinate on the surrounding accretion disk. The GRB is powered by the baryon poor outflows in the BZ process, and the associated SN is powered by very small fraction of the spin energy transferred from the BH to the disk in the magnetic coupling (MC) process. The timescale of the GRB is fitted by the duration of the open magnetic flux on the horizon. It turns out that the data of several GRB-SNe are well fitted with our model.

Subject headings: black hole physics–accretion disks–gamma-ray bursts–supernovae

1. INTRODUCTION

As is well known, the Blandford-Znajek (BZ) process is an effective mechanism for powering jets from quasars and active galactic nuclei (Blandford & Znajek 1977; Rees 1984). In the BZ process energy and angular momentum are extracted from a rotating black hole (BH), and transferred to the remote astrophysics load by the open magnetic field lines. Recently, much attention has been paid on the issue of the long duration gamma-ray bursts (GRBs) powered by the BZ process (Paczynski 1993, 1998; Meszaros & Rees 1997). A detailed model of GRBs invoking the BZ process is given by Lee, Wijers, & Brown 2000, hereafter L00), and they concluded that a fast-spinning BH with strong magnetic field $\sim 10^{15}G$ can provide energy $\sim 10^{53}ergs$ to power a GRB within 1000 seconds.

Recently the observations and theoretical considerations have linked long duration GRB with ultra-bright type Ib/c supernovae (Galama et al. 1998; Bloom et al. 1999; Galama et al. 2000). The first candidate was provided by SN1998bw and GRB980425, and the recent HETE-II burst GRB030329 has greatly enhanced the confidence in this association (Stanek et al. 2003; Hjorth et al. 2003).

¹Send offprint requests to: Ding-Xiong Wang (dxwang@hust.edu.cn)

Not long ago Brown et al. (2000, hereafter B00) worked out a specific scenario for GRB-SN connection. They argued that the GRB is powered by the BZ process, and the SN is powered by the energy dissipated into the disk through closed magnetic field lines coupling the disk with the BH. The latter energy mechanism is referred to as the magnetic coupling (MC) process, which is regarded as one of the variants of the BZ process (Blandford 1999; Putten 1999; Li 2000, 2002; Wang, Xiao & Lei 2002, hereafter W02). It is shown in B00 that about 10^{53}ergs are available to power both GRBs and SNe. However, they failed to distinguish the fractions of the energy for these two objects.

More recently, van Putten et al. (van Putten 2001a; van Putten & Levinson 2003; van Putten et al. 2004) worked out a poloidal topology for the open and closed magnetic field lines, where the separatrix on the horizon is defined by a finite half-opening angle. The duration of GRB is set by the lifetime of rapid spin of the BH. It is found that GRB and SN are powered by a small fraction of the BH spin energy. This result is consistent with the observations, i.e., the duration of GRB with tens of seconds, the true GRB energies distributed around $5 \times 10^{50} \text{ergs}$ (Frail et al. 2002), and the aspherical SNe kinetic energies with $2 \times 10^{51} \text{ergs}$ (Hoflich et al. 1999).

In this paper we propose a toy model for GRB-SN by considering coexistence of the BZ and MC processes (CEBZMC), where the configuration of the magnetic field is based the works of Wang et al. and van Putten et al. (W02; Wang et al. 2003, hereafter W03; van Putten 2001a; van Putten et al. 2004). The transferred energy and the duration for GRBs are calculated in the evolving process of the half-opening angle of the magnetic flux on the BH horizon, while the transferred energy for SNe is calculated in the MC process. Thus the association of GRBs and SNe is explained reasonably and consistently.

This paper is organized as follows. In § 2 the configuration of the magnetic field is described with the half-opening angle θ_{BZ} of the magnetic flux tube on the horizon, and we find that θ_{BZ} can evolve to zero in some range of the power-law index n of the magnetic field on the disk. In § 3 the toy model for GRB-SN based on BH evolution is described in the parameter space consisting of the BH spin a_* and the power-law index n . It is shown that the half-opening angle θ_{BZ} plays an important role of determining (i) the powers for GRBs and SNe, (ii) the duration of GRB. In § 4 we calculate the duration of GRB and the energies of GRB-SN, and compare with the other models of GRBs invoking the BZ process. It is found the duration of GRB obtained in our model is shorter than that of the other models. It turns out that this model is in excellent agreement with the observations, owing to considering the MC effects, while too much energy than the needed energy of GRB is produced in the other models without the MC effects. Finally, in § 5, we summarize our main results and discuss some problems concerning our model. Throughout this paper, the geometric units $G = c = 1$ are used.

2. CONFIGURATION OF THE MAGNETIC FIELD

Recently, we proposed a model of CEBZMC, in which the remote astrophysical load in the BZ process and the disk load in the MC process are connected with a rotating BH by open and closed magnetic field lines, respectively (W02; W03). The poloidal configuration of the magnetic field is shown in Figure 1, which is adapted from van Putten (2001a).

In Figure 1 the angle θ_{BZ} is the half-opening angle of the open magnetic flux tube, indicating the angular boundary between open and closed field lines on the horizon. This angle has been discussed by van Putten and his collaborators, who related the half-opening angle to the curvature in poloidal

topology of the inner torus magnetosphere (van Putten & Levinson 2003; van Putten et al. 2004). In W03 the angle θ_{BZ} is determined based on the following assumptions.

- (1) The theory of a stationary, axisymmetric magnetosphere formulated in the work of MacDonald & Thorne (1982) is applicable not only to the BZ process but also to the MC process. The magnetosphere is assumed to be force-free outside the BH and disk.
- (2) The disk is both stable and perfectly conducting, and the closed magnetic field lines are frozen in the disk. The disk is thin and Keplerian, and it lies in the equatorial plane of the BH, with inner boundary being at the marginally stable orbit.
- (3) The poloidal magnetic field is assumed to be constant on the horizon and to vary as a power law on the disk as follows (Blandford 1976),

$$B_D \propto \xi^{-n}, \quad (1)$$

where B_D is the magnetic field on the disk, the parameter n is the power-law index, and $\xi \equiv r/r_{ms}$ is the radial coordinate on the disk, which is defined in terms of the radius $r_{ms} \equiv M\chi_{ms}^2$ of the marginally stable orbit (Novikov & Thorne 1973).

- (4) The magnetic flux connecting the BH with its surrounding disk takes precedence over that connecting the BH with the remote load.

Assumption (4) is proposed based on two reasons: (i) the magnetic field on the horizon is brought and held by the surrounding magnetized disk, and (ii) the disk is much nearer to the BH than the remote load.

A mapping relation between the parameter ξ and the angular θ is derived in W03 based on the conservation of magnetic flux of the closed field lines.

$$\cos \theta = \int_1^\xi G(a_*; \xi, n) d\xi, \quad (2)$$

where $a_* \equiv J/M^2$ is the BH spin defined by the BH mass M and angular momentum J , and the function $G(a_*; \xi, n)$ is given by

$$G(a_*; \xi, n) = \frac{\xi^{1-n} \chi_{ms}^2 \sqrt{1 + a_*^2 \chi_{ms}^{-4} \xi^{-2} + 2a_*^2 \chi_{ms}^{-6} \xi^{-3}}}{2\sqrt{(1 + a_*^2 \chi_{ms}^{-4} + 2a_*^2 \chi_{ms}^{-6})(1 - 2\chi_{ms}^{-2} \xi^{-1} + a_*^2 \chi_{ms}^{-4} \xi^{-2})}}. \quad (3)$$

Based on the assumption (4) the angle θ_{BZ} can be determined by taking $\xi = \infty$ for the highest closed field lines as follows,

$$\cos \theta_{BZ} = \int_1^\infty G(a_*; \xi, n) d\xi. \quad (4)$$

The curves of θ_{BZ} versus a_* with different values of power-law index n are shown in Figure 2.

Inspecting Figure 2, we have the following results.

(1) The half-opening angle θ_{BZ} always increases monotonically with the increasing n .

(2) For some values of n , such as $n = 3.5, 4, 4.5, 5$ and 5.493 , the angle θ_{BZ} can evolve to zero with the decreasing a_* .

The second point implies that the open magnetic flux tube will be shut off when the BH spin decreases to the critical value a_*^{GRB} corresponding to $\theta_{BZ} = 0$. The lifetime of the half-opening angle θ_{BZ} is defined as the evolution time of BH from the initial spin $a_*(0)$ to a_*^{GRB} . Obviously it is less than the lifetime of BH spin, which is the evolution time of BH from $a_*(0)$ to zero. By using equation (4) we find that a_*^{GRB} exists only for a specific value range of the power-law index n , i.e., $3.003 \leq n \leq 5.493$. The half-opening angle θ_{BZ} remains zero for $n < 3.003$ in the whole evolving process, and it will never evolve to zero for $n > 5.493$.

3. A TOY MODEL FOR GRB-SN BASED ON BH EVOLUTION IN CEBZMC

Considering that the angular momentum is transferred from a rapidly rotating BH to the disk, on which a positive torque exerts, we think the accretion onto the BH might be probably halted. This state is essentially the suspended accretion state proposed by van Putten & Ostriker (2001), and the evolution of the BH is governed by the BZ and the MC processes.

The expressions for the BZ and MC powers and torques are derived in W02 by using an improved equivalent circuit for the BH magnetosphere based on the work of MacDonald and Thorne (1982). Considering the angular boundary θ_{BZ} , we express the BZ and MC powers and torques as follows.

$$\tilde{P}_{BZ} \equiv P_{BZ}/P_0 = 2a_*^2 \int_0^{\theta_{BZ}} \frac{k(1-k)\sin^3\theta d\theta}{2-(1-q)\sin^2\theta}, \quad (5)$$

$$\tilde{T}_{BZ} \equiv T_{BZ}/T_0 = 4a_*(1+q) \int_0^{\theta_{BZ}} \frac{(1-k)\sin^3\theta d\theta}{2-(1-q)\sin^2\theta}, \quad (6)$$

$$\tilde{P}_{MC} \equiv P_{MC}/P_0 = 2a_*^2 \int_{\theta_{BZ}}^{\pi/2} \frac{\beta(1-\beta)\sin^3\theta d\theta}{2-(1-q)\sin^2\theta}, \quad (7)$$

$$\tilde{T}_{MC} \equiv T_{MC}/T_0 = 4a_*(1+q) \int_{\theta_{BZ}}^{\pi/2} \frac{(1-\beta)\sin^3\theta d\theta}{2-(1-q)\sin^2\theta}, \quad (8)$$

where we have $q \equiv \sqrt{1-a_*^2}$ and

$$\begin{cases} P_0 \equiv \langle B_H^2 \rangle M^2 \approx 6.59 \times 10^{50} \times \left(\frac{B_H}{10^{15}G}\right)^2 \left(\frac{M}{M_\odot}\right)^2 \text{ erg} \cdot s^{-1} \\ T_0 \equiv \langle B_H^2 \rangle M^3 \approx 3.26 \times 10^{45} \left(\frac{B_H}{10^{15}G}\right)^2 \left(\frac{M}{M_\odot}\right)^3 g \cdot cm^2 \cdot s^{-2} \end{cases} \quad (9)$$

In equation (9) B_H is the magnetic field on the BH horizon. The parameters k and β are the ratios of the angular velocities of the open and closed magnetic field lines to that of the BH, respectively.

Usually, $k = 0.5$ is taken for the optimal BZ power. Since the closed field lines are assumed to be frozen in the disk, the ratio β is related to a_* and ξ by

$$\beta \equiv \Omega_D/\Omega_H = \frac{2(1+q)}{a_* \left[(\sqrt{\xi}\chi_{ms})^3 + a_* \right]}, \quad (10)$$

where Ω_H and Ω_D are the angular velocities of the BH and the disk, respectively.

Since θ_{BZ} in equations (5) and (7) is determined by a_* and n , we have the curves of the powers $\tilde{P}_{BZ}$ and $\tilde{P}_{MC}$ versus a_* for the given values of n as shown in Figure 3.

Inspecting Figure 3, we have the following results.

(1) The MC power P_{MC} is always greater than the BZ power P_{BZ} for a wide value range of the power-law index n , provided that the BH spin a_* is not very small.

(2) For the given values of a_* the BZ power P_{BZ} increases with the increasing n , while the MC power P_{MC} decreases with it.

(3) The power $P_{MC} = 0$ holds at some small values of a_* for the given values of the index n .

From equation (5) we find $P_{BZ} = 0$ when $\theta_{BZ} = 0$. Therefore the effective time of the BZ process can be set by the lifetime of θ_{BZ} , i.e., the evolution time of BH from $a_*(0)$ to a_*^{GRB} for $3.003 \leq n \leq 5.493$. As argued in our previous work (Wang, Lei & Ma 2003), we have $P_{MC} = 0$ when the energy transported from the BH to the disk is equal to that from the disk to the BH. Henceforth the corresponding critical BH spin is denoted by a_*^{SN} . We have $P_{MC} > 0$ for $a_* > a_*^{SN}$, and $P_{MC} < 0$ for $a_* < a_*^{SN}$, indicating that the energy transferred from the BH to the disk dominates that transferred in the inverse direction.

Based on the conservation laws of energy and angular momentum we have the following evolution equations of the rotating BH,

$$dM/dt = -(P_{BZ} + P_{MC}), \quad (11)$$

$$dJ/dt = -(T_{BZ} + T_{MC}). \quad (12)$$

Incorporating equations (11) and (12), we have the evolution equation for the BH spin expressed by

$$da_*/dt = -M^{-2}(T_{BZ} + T_{MC}) + 2M^{-1}a_*(P_{BZ} + P_{MC}) = B_H^2 MA(a_*, n) \quad (13)$$

where the function $A(a_*, n)$ is

$$A(a_*, n) = 2a_*\tilde{P}_{mag} - \tilde{T}_{mag}, \quad (14)$$

where

$$\tilde{P}_{mag} = \tilde{P}_{BZ} + \tilde{P}_{MC}, \quad \tilde{T}_{mag} = \tilde{T}_{BZ} + \tilde{T}_{MC}. \quad (15)$$

The sign of $A(a_*, n)$ determines whether a_* decreases or increases, and we have the curves of $A(a_*, n)$ versus a_* for different values of n as shown in Figure 4.

From Figure 4, we find the following evolution characteristics of the BH spin.

(1) The function $A(a_*, n) > 0$ for $a_* < a_*^{eq}$, and $A(a_*, n) < 0$ for $a_* > a_*^{eq}$, and the BH spin always evolve to the equilibrium spin a_*^{eq} .

(2) Both the equilibrium spin a_*^{eq} and the decreasing rate of the BH spin are sensitive to the power-law index n . The greater the index n is, the greater is the equilibrium spin a_*^{eq} , and the more slowly the BH spin decreases to a_*^{eq} .

(3) The features of our model for GRB-SN can be described by three functions: $A(a_*, n)$, $\tilde{P}_{BZ}(a_*, n)$ and $\tilde{P}_{MC}(a_*, n)$.

Setting $A(a_*, n) = 0$, $\tilde{P}_{BZ}(a_*, n) = 0$ and $\tilde{P}_{MC}(a_*, n) = 0$, we have the three characteristic curves in the parameter space as shown in Figure 5.

The parameter space is divided into seven regions, **IA**, **IB**, **II**, **IIIA**, **IIIB**, **IVA** and **IVB**, where the points **M** and **N** are the intersections of the curve $\tilde{P}_{BZ}(a_*, n) = 0$ with the curves $\tilde{P}_{MC}(a_*, n) = 0$ and $A(a_*, n) = 0$, respectively. **IA** and **IB** is divided by the dashed line starting from the point **M**. By using the equations (5), (7) and (14) we obtain the coordinates of points **M** and **N**, i.e., **M**(0.253, 5.067) and **N**(0.222, 5.122), respectively. In Figure 5 each black dot with an arrowhead in these sub-regions is referred to as a representative point (**RP**), which represents one evolution state of the BH as shown in Table 1.

It is found from Figure 5 and Table 1 that GRB-SN only occurs in region **I** and **I** corresponding to CEBZMC. Based on the observations that GRBs need less energy than SNe (B00), we think that the evolving path of the **RP** in region **IA** is reasonable, which corresponds to $3.003 < n < 5.067$, and the RP attains $\tilde{P}_{BZ}(a_*, n) = 0$ with $\tilde{P}_{MC}(a_*, n) > 0$ in braking the spinning BH.

Based on the above discussion we describe the toy model for GRB-SN in the parameter space as follows.

(1) The fast-spinning BH can power GRBs and SNe via the BZ and MC processes, respectively, which is represented by the **RP** in region **IA**. The GRB will be shut off, when the **RP** reaches eventually the curve $\tilde{P}_{BZ}(a_*, n) = 0$.

(2) Energy is transferred from the spinning BH to the disk via the MC process in the evolving path from region **IA** to region **II**. The SN will be shut off with no further energy transferred into the disk, if the **RP** arrives at the curve $\tilde{P}_{MC}(a_*, n) = 0$.

(3) Since enormous energy is deposited in the disk due to the MC process, the disk might be destroyed eventually, leaving the BH alone without the disk. This outcome probably arises from explosion of SNe (B00).

4. ENERGIES AND TIME SCALES FOR GRB-SN

In the evolving path of the **RP** going through regions **IA**, **II** and **IIIA** we have $a_*^{GRB} \geq a_*^{SN} > a_*^{eq}$. Based on the above discussion on correlation of the BH evolution with the association of GRB-SN, the energy E_{BZ} and E_{SN} extracted in the BZ and MC processes are given respectively by

$$E_{BZ} = \int_{a_*(0)}^{a_*^{GRB}} \frac{P_{BZ}}{(da_*/dt)} da_* = 1.79 \times 10^{54} \text{ergs} \times \left(\frac{M(0)}{M_\odot} \right) \int_{a_*(0)}^{a_*^{GRB}} \frac{\tilde{M} \tilde{P}_{BZ}}{-\tilde{T}_{mag} + 2a_* \tilde{P}_{mag}} da_*, \quad (16)$$

$$E_{MC} = \int_{a_*(0)}^{a_*^{SN}} \frac{P_{MC}}{(da_*/dt)} da_* = 1.79 \times 10^{54} \text{ergs} \times \left(\frac{M(0)}{M_\odot} \right) \int_{a_*(0)}^{a_*^{SN}} \frac{\tilde{M} \tilde{P}_{MC}}{-\tilde{T}_{mag} + 2a_* \tilde{P}_{mag}} da_*. \quad (17)$$

The true energy for GRBs, E_γ , and the energy for SNe, E_{SN} , are related respectively to E_{BZ} and E_{MC} by

$$E_\gamma = \varepsilon_\gamma E_{BZ}, \quad (18)$$

$$E_{SN} = \varepsilon_{SN} E_{MC}, \quad (19)$$

where ε_γ and ε_{SN} denote the efficiencies of converting E_{BZ} and E_{MC} into E_γ and E_{SN} , respectively. Following van Putten et al. (2004), we take $\varepsilon_\gamma = 0.15$ in calculations. The duration of GRB, t_{GRB} , is defined as the lifetime of the half-opening angle θ_{BZ} , which is exactly equal to the time for the BH spin evolving from $a_*(0)$ to a_*^{GRB} , i.e.,

$$t_{GRB} = 2.7 \times 10^3 \text{s} \times \left(\frac{10^{15} G}{B_H} \right)^2 \left(\frac{M_\odot}{M(0)} \right) \int_{a_*(0)}^{a_*^{GRB}} \frac{\tilde{M}^{-1}}{-\tilde{T}_{mag} + 2a_* \tilde{P}_{mag}} da_*, \quad (20)$$

where $\tilde{M}$ is the ratio of the BH mass to its initial value $M(0)$, and it is calculated by

$$\tilde{M} \equiv \frac{M}{M(0)} = \exp \int_{a_*(0)}^{a_*} \frac{da_*}{\tilde{T}_{mag} / \tilde{P}_{mag} - 2a_*}. \quad (21)$$

Equation (21) can be derived by using equations (11) and (13). Obviously, t_{GRB} is shorter than the lifetime of the BH spin evolving from $a_*(0)$ to $a_* = 0$.

Considering that spin-energy of the BHs produced in core-collapse is around 50% of maximum or less in centered nucleation (van Putten 2004), we take the initial BH spin as $a_*(0) = 0.9$ in equations (16), (17) and (21). The value range $3.534 < n < 5.067$ is taken in the evolving path of the **RP** in region **IA** corresponding to GRB-SN. In addition, $M(0) = 7M_\odot$ and $B_H = 10^{15} G$ are assumed, and the cutoff of T_{GRB} is taken as T_{90} in calculations, which is the duration for 90% of the total BZ energy to be extracted (Lee & Kim 2002).

As is well know, the rotational energy E_{rot} of a Kerr BH is expressed by (Thorne, Price & Macdonald 1986)

$$E_{rot} = f(a_*)M(0), \quad (22)$$

where

$$f(a_*) = 1 - \sqrt{\frac{1}{2}(1 + \sqrt{1 - a_*^2})}. \quad (23)$$

Substituting $a_*(0) = 0.9$ and $M(0) = 7M_\odot$ into equation (22), we have the total rotational energy of the BH,

$$E_{rot} \approx 1.9 \times 10^{54} \text{ergs}. \quad (24)$$

To compare rotational energy of the BH extracted in the BZ and MC processes, we calculate the ratios of E_{BZ} and E_{MC} to E_{rot} , i.e.,

$$\eta_{BZ} \equiv E_{BZ}/E_{rot}, \quad \eta_{MC} \equiv E_{MC}/E_{rot}. \quad (25)$$

The curves of η_{BZ} and η_{MC} versus n are shown in Figure 6.

From Figure 6 we find that E_{MC} is significantly greater than E_{BZ} , and the greater the index n , the more energy extracted by BZ process. These results can be also obtained by inspecting Figures 3 and 5. The ratio η_{MC} varies from 25.8% to 33.8% with the ratio η_{BZ} varying from zero to 3.7% for $3.534 < n < 5.067$. Combining equations (24) and (25) with the above ratios, we obtain E_{MC} varying from $4.929 \times 10^{53} \text{ergs}$ to $6.459 \times 10^{53} \text{ergs}$ with $E_{BZ} = 7.003 \times 10^{52} \text{ergs}$. And the energy for GRB is inferred by using $\varepsilon_\gamma = 0.15$.

Calculations of SN1998bw with aspherical geometry show that the needed kinetic energy is about $2 \times 10^{51} \text{ergs}$ (Hoflich et al. 1999), and this results in the ratio ε_{SN} varying from 0.003 to 0.004 for E_{MC} varying from $6.459 \times 10^{53} \text{ergs}$ to $4.929 \times 10^{53} \text{ergs}$ in equation (19). In this paper we take the average value, $\varepsilon_{SN} \approx 0.0035$.

It has been shown that most fraction of the rotational energy of a Kerr BH is emitted in unseen channels, such as in gravitational radiation and MeV-neutrino emissions (van Putten 2001a, 2001b; van Putten and Levinson 2002). The above efficiency factors obtained in our model, $\varepsilon_\gamma = 0.15$ and $\varepsilon_{SN} \approx 0.0035$, imply that only very small fraction of E_{rot} is converted into the energy for GRBs and SNe. These results are consistent with those obtained by van Putten and his collaborators.

By using equation (20) the duration of GRBs, T_{90} , versus the parameter n is shown in Figure 7.

We find that the estimated duration of GRB is tens of seconds, which is consistent with the observations (Kouveliotous et al. 1993). For $n = 4$ and $B_H = 1 \times 10^{15} G$ in the evolving path of the **RP** from $a_*(0) = 0.9$ to $a_*^{SN} = 0.235$, we obtain the duration of GRB $T_{90} = 21s$, and the energies for GRB and SN are $E_\gamma = 5 \times 10^{50} \text{ergs}(\frac{\varepsilon_\gamma}{0.15})$ and $E_{SN} = 2 \times 10^{51} \text{ergs}(\frac{\varepsilon_{SN}}{0.0035})$, respectively. These results are in excellent agreement with the observed durations of tens of seconds (Kouveliotous et al. 1993), energies

$E_\gamma = 5 \times 10^{50} \text{ergs}$ in gamma rays (Frail et al. 2002), and inferred kinetic energy $E_{SN} = 2 \times 10^{51} \text{ergs}$ in SN1998bw with aspherical geometry (Hoflich et al. 1999).

For several GRB-SNe, the observed energy E_γ and duration T_{90} can be fitted by adjusting the parameters n and B_H , and the energy E_{SN} can be predicted as shown in Table 2.

In the previous model for GRBs powered by the BZ process, the duration of GRB is estimated either in the case that the BH is spun down to zero or in the case that the whole disk is plunged into the BH (L00; Lee & Kim 2000, 2002; Wang, Lei & Xiao 2002). However, the MC effects have not been taken into account in these models. In this model the duration of GRB is estimated by the lifetime of the half-opening angle on the BH horizon based on CEBZMC. It turns out that both the BZ and MC processes play very important roles for GRN-SN in this model.

The quantities, E_γ and T_{90} , calculated in the three different models for GRBs are listed in Table 3, where the abbreviations **CEBZMC**, **BZO** and **BZACC** represent this model, the model invoking the BZ process only, the model invoking the BZ process with disk accretion, respectively.

It is found in Table 3 that the energy and duration time obtained in this model are in excellent agreement with the observed durations of tens of seconds and energies $E_\gamma \approx 5 \times 10^{50} \text{ergs}$. However the energies obtained in the models **BZO** and **BZACC** seem too much to fit the observations.

5. DISCUSSION

In this paper we discuss a toy model for GRB-SN, where the energy for GRBs and SNe are powered by the BZ and MC processes, respectively, and the duration for GRB is estimated by the lifetime of the half-opening angle θ_{BZ} . The energy extracted in the BZ process for GRBs is much less than that extracted in the MC process for SNe. This result is consistent with the observations: the true energy of GRB $E_\gamma = 5 \times 10^{50} \text{ergs}$ and the aspherical SNe of kinetic energy $E_{SN} = 2 \times 10^{51} \text{ergs}$. For a set of GRBs the observed true energy E_γ and duration T_{90} can be well fitted by taking adequate values of the power-law index n and magnetic field B_H . However, there are still several issues related to this model.

(i) In this paper we take the evolving path of the **RP** in the region **IA** to describe the association of GRB-SN as shown in Figure 5. In this case the GRB will terminate before the SN event stops. In fact the evolving path in region **IB** can be also used to describe the association of GRB-SN, and it implies that the **RP** will arrive at the characteristic curve of $\tilde{P}_{MC} = 0$ ahead of reaching the characteristic curve of $\tilde{P}_{BZ} = 0$. However, we think that the GRB will still terminate before the SN event stops, because the disk will be destroyed during the explosion of the SN, and the BZ process can not work without the magnetic field supported by its surrounding disk.

(ii) In order to highlight the effects of the BZ and MC processes in powering GRB-SN, we neglect the effects of the disk accretion in this model. In fact, disk accretion plays a very important role in BH evolution. The slowing down of the BH spin will be delayed significantly since the accreting matter delivers remarkable amount of energy and angular momentum to the BH. Strictly speaking, disk accretion cannot be halted steadily by the MC process. The MC effects on disk accretion is a very complicated, and it should be treated dynamically.

(iii) According to Kruskal-Shafranov criteria (Kadomtsev 1966), the screw instability will occur if the toroidal magnetic field becomes so strong that the magnetic field lines turns around itself about once.

In our latest work (Wang et al. 2004), we argued that the state of CEBZMC always accompanies the screw instability. In this model the half-opening angle θ_{BZ} is related to the infinite radial parameter ξ by equation (4). We expect to derive a greater θ_{BZ} by considering the restriction of the screw instability to the MC region. Thus the ratios η_{GRB} and η_{SN} will be changed. We shall improve this model in future.

6. Acknowledgments

We thank the anonymous referees for numerous constructive suggestions. This work is supported by the National Natural Science Foundation of China under grants 10173004, 10373006 and 10121503.

REFERENCES

- Blandford, R. D. 1976, MNRAS, 176, 465
- Blandford, R. D. 1999, in ASP Conf. Ser. 160, Astrophysical Discs: An EC Summer School, ed. J. A. Sellwood & J. Goodman (San Francisco: ASP), 265
- Blandford, R. D., & Znajek, R. L. 1977, MNRAS, 179, 433
- Bloom, J. S. et al. 1999, Nature, 401, 453
- Brown, G. E., et al. 2000, New Astronomy 5, 191 (B00)
- Costa, E., et al. 1997, IAUC Circ. No. 6649
- Dar, A. 2004, astro-ph/0405386
- Frail, D. A., et al. 2002, ApJ, 562, L55
- Galama, T. J., et al. 1998, Nature, 395, 672
- Galama, T. J., et al. 2000, ApJ, 536, 185
- Heise, J., et al. 1999, IAUC Circ. No. 7221
- Hjorth, J., et al. 2003, Nature, 423, 847
- Hoflich, P. J., Wheeler, J. C., & Wang, L. 1999, ApJ, 521, 179
- Kadomtsev, B. B. 1966, Rev. Plasma Phys., 2, 153
- Kouveliotou et al. 1993, Astrophys. J. Lett., 413, L101
- Lee, H. K., & Wijers, R. A. M. J., Brown, G. E. 2000, Phys. Rep., 325, 83 (L00)
- Lee, H. K., & Kim, H. K. 2000, J. Korean Phys. Soc. 36, 188
- Lee, H. K., & Kim, H. K. 2002, J. Korean Phys. Soc. 40, 524

- Li, L. -X. 2000, ApJ, 533, L115
- Li, L. -X. 2002, ApJ, 567, 463
- MacDonald, D., & Thorne, K. S. 1982, MNRAS, 198, 345
- Meszaros P., & Rees M. J. 1997, ApJ, 482, L29
- Novikov, I. D., & Thorne, K. S. 1973, in Black Holes, ed. Dewitt C (Gordon and Breach, New York) p.345
- Paczynski, B. 1993, in Relativistic Astrophysics and Particle Cosmology, ed. C. W. Akerlof & M. A. Srednicki (Ann. NY Acad. Sci., Vol. 688), 321
- Paczynski, B. 1998, ApJ, 494, L45
- Rees, M. J. 1984, ARA&A. 22, 471
- Stanek, K. Z. et al. 2003, ApJ, 591, L17
- Thorne, K. S., Price, R. H., & Macdonald, D. A. 1986, *Black Holes: The Membrane Paradigm*, Yale Univ. Press, New Haven
- van Putten, M. H. P. M., 1999, Science, 284, 115
- van Putten, M. H. P. M. 2001a, Phys. Rep., 345, 1
- van Putten, M. H. P. M. 2001b, Phys. Rev. Lett., 87, 091101
- van Putten, M. H. P. M., & E. C. Ostriker, E. C. 2001, ApJ, 552, L31
- van Putten, M. H. P. M., Levinson, A. 2002, Science, 295, 1874
- van Putten, M. H. P. M., Levinson, A. 2003, ApJ, 584, 937
- van Putten, M. H. P. M., Levinson, A., Regimbau, T., Punturo, M., & Harry, G. M. 2004, Phys. Rev. D, 69, 044007
- van Putten, M. H. P. M. 2004, astro-ph/0407112 (to appear in ApJ Lett.)
- Wang, D.-X., Lei, W.-H., & Xiao, K. 2002, ApJ, 580, 358
- Wang, D.-X., Xiao, K., & Lei, W.-H. 2002, MNRAS, 335, 655 (W02)
- Wang, D.-X., Ma, R.-Y., Lei, W.-H., & Yao, G.-Z. 2003, ApJ, 595, 109 (W03)
- Wang, D.-X., Lei W.-H., & Ma, R.-Y. 2003, MNRAS, 342, 851
- Wang, D.-X., Ma, R.-Y., Lei, W.-H., & Yao, G.-Z. 2004, ApJ, 601, 1031

Table 1. Characteristic of BH evolution in each region of the parameter space

Region	da_*/dt	P_{BZ}	P_{MC}	RP displacement	Possible events
IA	< 0	> 0	> 0	Towards the left	GRB-SN
IB	< 0	> 0	> 0	Towards the left	GRB-SN
II	< 0	—	> 0	Towards the left	SN
IIIA	< 0	—	< 0	Towards the left	—
IIIB	< 0	> 0	< 0	Towards the left	GRB
IVA	> 0	—	< 0	Towards the right	—
IVB	> 0	> 0	< 0	Towards the right	GRB

Table 2. Four GRBs of the given true energy E_γ and T_{90} s fitted with the different power-law index n and the predicted E_{SN}

GRB ^a	$E_\gamma(10^{51}ergs)^b$	$T_{90}(s)$	n	$B_H(10^{15}G)$	E_{SN}^c
970508	0.234	15 ^d	3.885	0.97	$1.947 \times 10^{51}ergs(\frac{\epsilon_{SN}}{0.0035})$
990712	0.445	30 ^e	3.975	0.81	$1.989 \times 10^{51}ergs(\frac{\epsilon_{SN}}{0.0035})$
991208	0.455	39.84 ^f	3.985	0.75	$1.995 \times 10^{51}ergs(\frac{\epsilon_{SN}}{0.0035})$
991216	0.695	7.51 ^f	4.058	1.80	$2.032 \times 10^{51}ergs(\frac{\epsilon_{SN}}{0.0035})$

^aThe sample of GRB-SN takes from the sample listed in Table 1 of Dar (2004)

^bThe true energy E_γ of GRB refers to the sample listed in Table 1 of Frail et al (2002)

^cThe predicted energy of SNe based on our model

^dThe duration of GRB970508 refers to Costa et al (1997)

^eThe duration of GRB990712 refers to Heise et al. (1999)

^fThe durations of GRB991208 and GRB991216 refer to the sample listed in Table 2 of Lee & Kim (2002)

Table 3. Energy and duration obtained in three models for GRBs with $a_*(0) = 0.9$, $M_H(0) = 7M_\odot$, $B_H = 10^{15}G$ and the initial disk mass $M_D(0) = 3M_\odot$

Model	$E_\gamma(ergs)$	$T_{90}(s)$
BZO	1.225×10^{53}	348
BZACC	1.734×10^{53}	397
CEBZMC	$< 1.050 \times 10^{52}$	< 115




Fig. 1.— Poloidal magnetic field connecting a rotating BH with remote astrophysical load and a surrounding disk




Fig. 2.— Curves of θ_{BZ} vs. a_* for $n = 3.5, 4, 4.5, 5, 5.493$ and 6 .




Fig. 3.— Curves of $\tilde{P}_{BZ}$ (dotted lines) and $\tilde{P}_{MC}$ (solid lines) vs. a_* with $n=3.5, 4.5$ and 5.493 for $0 < a_* < 1$.




Fig. 4.— Curves of $A(a_*, n)$ vs. a_* for $n=3.5, 4, 4.5$ and 5.493 .




Fig. 5.— Parameter space of BH evolution in CEBZMC.




Fig. 6.— Curves of η_{BZ} (dashed line) and η_{MC} (solid line) vs. n for $3.534 < n < 5.067$.




Fig. 7.— Curve of T_{90} vs. n for $3.534 < n < 5.067$.